

Erratum

Quantum Cosmology With Conformally Invariant Scalar Field

(Physics Letters A 221 (1996) 359)

NIVALDO A. LEMOS

Departamento de Física

Universidade Federal Fluminense

Av. Litorânea s/n, Boa Viagem

24210-340, Niterói, RJ - Brazil

e-mail: nivaldo@if.uff.br

The paper contains a harmless typographical mistake and, regrettably, an algebraic error with serious consequences. The prefactor in the last term of Eq.(2) should be $1/6$ instead of 3 , and the last term in Eq.(7) should include k as a factor. As a consequence, in the case $k = \Lambda = 0$ considered in the paper Eq.(8) should read

$$p_R^2 - p_\chi^2 = 0 \quad .$$

In the second of Eqs.(9) the term $-4\chi^2$ within parentheses should be deleted, and the right-hand side of the last of Eqs.(9) should be zero. The classical equations of motion are solved by

$$R(t) = At \quad , \quad \chi(t) = \pm At + B \quad ,$$

instead of Eq.(10). In Section 3 one should replace Eqs.(12) and (13) by

$$H = -p_R = \sqrt{p_\chi^2} \quad , \quad \hat{H} = \sqrt{\hat{p}_\chi^2} \quad ,$$

whereas Eqs.(14) and (15) should read

$$\varphi_p(\chi) = e^{ip\chi} \quad , \quad \hat{H}\varphi_p(\chi) = |p|\varphi_p(\chi) \quad , \quad p \in \mathbf{R} \; .$$

The reduced-phase-space discussion of singularities in the gauge $t = R$ is still valid, and Eq.(23) remains true *for small* t if one just replaces $(n + 1/2)/2$ by $\langle \varphi | \hat{\chi}^2 | \varphi \rangle$ where $\varphi(\chi)$ is a square-integrable wave function at $t = 0$. The analysis based on the Wheeler-DeWitt equation, however, is now entirely different from the one given in the paper, which must be disregarded. Although Eq.(25) is to be replaced by

$$\hat{H} = -\sqrt{\hat{p}_R^2} \; ,$$

the subsequent conclusions in Section 5 remain unchanged.

Thanks are due to Nelson Pinto-Neto, José Acácio de Barros e Marco A. Sagiore-Leal for pointing out the errors.